

Noether identities of a generic differential operator. The Koszul–Tate complex

G. SARDANASHVILY

Department of Theoretical Physics, Moscow State University

Given a generic Lagrangian system, its Euler–Lagrange operator obeys Noether identities which need not be independent, but satisfy first-stage Noether identities, and so on. This construction is generalized to arbitrary differential operators on a smooth fiber bundle. Namely, if a certain necessary and sufficient condition holds, one can associate to a differential operator the exact chain complex with the boundary operator whose nilpotency restarts all the Noether identities characterizing the degeneracy of an original differential operator.

1 Introduction

Given a generic Lagrangian system on a fiber bundle, its Euler–Lagrange operator $\mathcal{E}$ obeys the Noether identities. They need not be independent, but satisfy the first-stage Noether identities, which in turn are subject to the second-order Noether ones, and so on. The hierarchy of these Noether identities characterizes the degeneracy of a Lagrangian system in full. Noether’s second theorem states the relation between Noether identities and gauge symmetries of a Lagrangian [1, 2], but Noether identities can be introduced without regarding $\mathcal{E}$ as an Euler–Lagrange operator. Therefore, one can extend the notion of Noether identities to a generic differential operator on a fiber bundle. Our goal is the following.

Let $Y \rightarrow X$ be a smooth fiber bundle and $J^r Y$, $r = 1, \dots$, the jet manifolds of its sections (the index $r = 0$ further stands for Y). Let $E \rightarrow X$ be a vector bundle. A E -valued r -order differential operator $\mathcal{E}$ on Y is conventionally defined as a bundle morphism of $J^r Y$ to E over X [3, 4]. We associate to $\mathcal{E}$ the chain complex (10) whose boundaries vanish on $\text{Ker } \mathcal{E}$ (Proposition 2). It is a complex of a certain ring $\mathcal{S}_\infty[E; Y]$ of Grassmann-graded functions and their jets on the infinite order jet manifold $J^\infty Y$ of Y . For our purpose, this complex can be replaced with the short zero-exact complex $\mathcal{S}_\infty[E; Y]_{\leq 2}$ (11). Recall that a chain complex is called r -exact if its homology of degree $k \leq r$ is trivial.

Noether identities of a differential operator $\mathcal{E}$ are defined as nontrivial elements of the first homology $H_1(\delta)$ of the complex (11) (Definition 3). If this homology is finitely generated, the complex (11) can be extended to the one exact-complex $\mathcal{P}_\infty[E; Y; E_0]_{\leq 3}$ (19) with the boundary operator δ_0 (18) whose nilpotency conditions are equivalent to the above-mentioned Noether identities (Proposition 4). First-stage Noether identities are defined as two-cycles of this complex. They are trivial if two-cycles are boundaries, but the converse need not be true. Trivial first-stage Noether identities are boundaries iff a certain homology condition (called the two-homology regularity condition) holds (Proposition 6). In this case, the first-stage Noether identities are identified to nontrivial elements of the

second homology of the complex (19). If this homology is finitely generated, the complex (19) is extended to the two-exact complex $\mathcal{P}_\infty[E_1 E; Y; E_0]_{\leq 4}$ (30) with the boundary operator δ_1 (29) whose nilpotency conditions are equivalent to the Noether and first-stage Noether identities (Proposition 8). If the third homology of this complex is not trivial, the second-stage Noether identities exist, and so on. Iterating the arguments, we come to the following.

Let us have the $(N + 1)$ -exact complex $\mathcal{P}_\infty\{N\}_{\leq N+3}$ (33) such that: (i) the nilpotency conditions of its boundary operator δ_N (31) reproduce Noether and k -stage Noether identities for $k \leq N$, (ii) the $(N + 1)$ -homology regularity condition (Definition 9) holds. Then the $(N + 1)$ -stage Noether identities are defined as $(N + 2)$ -cycles of this complex. They are trivial if cycles are boundaries, while the converse is true iff the $(N + 2)$ -homology regularity condition is satisfied. In this case, $(N + 1)$ -stage Noether identities are identified to nontrivial elements of the $(N + 2)$ -homology of the complex (33) (item (i) of Theorem 11). If this homology is finitely generated, this complex is extended to the $(N + 2)$ -exact complex $\mathcal{P}_\infty\{N + 1\}_{\leq N+4}$ (41) with the boundary operator δ_{N+1} (40) whose nilpotency restarts all the Noether identities up to stage $(N + 1)$ (item (ii) of Theorem 11).

This iteration procedure results in the exact Koszul–Tate complex with the boundary operator whose nilpotency conditions reproduce all Noether and higher Noether identities characterizing the degeneracy of a differential operator $\mathcal{E}$

2 The ring of Grassmann-graded functions and their jets

All chain complexes considered in the article are complexes of certain rings of Grassmann-graded function and their jets on the infinite order jet manifold $J^\infty Y$ of a fiber bundle $Y \rightarrow X$. Let us describe such a ring.

Recall that $J^\infty Y$, is the projective limit $(\pi_r^\infty : J^\infty Y \rightarrow J^r Y)$ of the inverse system of jet manifolds

$$X \xleftarrow{\pi} Y \xleftarrow{\pi_0^1} J^1 Y \xleftarrow{\dots} J^{r-1} Y \xleftarrow{\pi_{r-1}^r} J^r Y \xleftarrow{\dots}, \quad (1)$$

of $J^\infty Y$, where π_{r-1}^r are affine bundles. It is a Fréchet manifold. A bundle atlas $\{(U_Y; x^\lambda, y^i)\}$ of $Y \rightarrow X$ induces the coordinate atlas

$$\{((\pi_0^\infty)^{-1}(U_Y); x^\lambda, y_\Lambda^i)\}, \quad y_{\lambda+\Lambda}^i = \frac{\partial x^\mu}{\partial x'^\lambda} d_\mu y_\Lambda^i, \quad 0 \leq |\Lambda|, \quad (2)$$

where $\Lambda = (\lambda_1 \dots \lambda_k)$, $\lambda + \Lambda = (\lambda \lambda_1 \dots \lambda_k)$ are symmetric multi-indices and

$$d_\lambda = \partial_\lambda + \sum_{0 \leq |\Lambda|} y_{\lambda+\Lambda}^i \partial_i^\Lambda, \quad d_\Lambda = d_{\lambda_1} \circ \dots \circ d_{\lambda_k},$$

are total derivatives. We further assume that the cover $\{\pi(U_Y)\}$ of X is also the cover of atlases of all vector bundles over X in question. The inverse system (1) yields the direct system

$$C^\infty(X) \xrightarrow{\pi^*} C^\infty(Y) \xrightarrow{\pi_0^{1*}} C^\infty(J^1 Y) \longrightarrow \dots C^\infty(J^{r-1} Y) \xrightarrow{\pi_{r-1}^{r*}} \dots$$

of rings of smooth real functions on jet manifolds $J^r Y$ with respect to the pull-back monomorphisms π_{r-1}^* . Its direct limit is the ring $\mathcal{O}_\infty Y$ of all smooth real functions on finite order jet manifolds modulo the pull-back identification.

Let us extend the ring $\mathcal{O}_\infty Y$ to a ring of graded functions on graded manifolds whose bodies are jet manifolds $J^r Y$ of Y [2, 5]. We restrict our consideration to graded manifolds $(Z, \mathfrak{A})$ with structure sheaves $\mathfrak{A}$ of Grassmann algebras of finite rank, and refer to the following Serre–Swan theorem.

Theorem 1. *Let Z be a smooth manifold. A Grassmann algebra $\mathcal{A}$ over the ring $C^\infty(Z)$ of smooth real functions on Z is isomorphic to the Grassmann algebra of graded functions on a graded manifold with a body Z iff it is the exterior algebra of some projective $C^\infty(Z)$ -module of finite rank.*

Proof. The proof follows at once from the Batchelor theorem [6] and the Serre–Swan theorem generalized to an arbitrary smooth manifold [7, 8]. The Batchelor theorem states that any graded manifold $(Z, \mathfrak{A})$ with a body Z is isomorphic to the one $(Z, \mathfrak{A}_Q)$ with the structure sheaf $\mathfrak{A}_Q$ of germs of sections of the exterior bundle

$$\wedge Q^* = \mathbb{R} \oplus_Z Q^* \oplus_Z \wedge^2 Q^* \oplus \cdots,$$

where Q^* is the dual of some vector bundle $Q \rightarrow Z$. We agree to call $(Z, \mathfrak{A}_Q)$ the simple graded manifold with the structure vector bundle Q . Its structure ring $\mathcal{A}_Q$ of graded functions (sections of $\mathfrak{A}_Q$) is the $\mathbb{Z}_2$ -graded exterior algebra of the $C^\infty(Z)$ -module of sections of $Q^* \rightarrow Z$. By virtue of the Serre–Swan theorem, a $C^\infty(Z)$ -module is isomorphic to the module of sections of a smooth vector bundle over Z iff it is a projective module of finite rank. $\square$

Remark 1. With respect to bundle coordinates (z^A, q^a) on Q and the corresponding fiber basis $\{c^a\}$ for $Q^* \rightarrow X$, graded functions read

$$f = \sum_{k=0} \frac{1}{k!} f_{a_1 \dots a_k} c^{a_1} \cdots c^{a_k}, \quad f \in C^\infty(Z), \quad (3)$$

where we omit the symbol of the exterior product of elements c^a . Let $u \in \mathfrak{d}\mathcal{A}_Q$ be a graded derivation of the $\mathbb{R}$ -ring $\mathcal{A}_Q$. Due to the canonical splitting $VQ = Q \times Q$, the fiber basis $\{\partial_a\}$ for vertical tangent bundle $VQ \rightarrow Q$ of $Q \rightarrow Z$ is the dual of $\{c^a\}$ and, therefore, u takes the local form $u = u^A \partial_A + u^a \partial_a$, where u^A, u^a are local graded functions [6]. It acts on graded functions (3) by the rule

$$u(f_{a \dots b} c^a \cdots c^b) = u^A \partial_A (f_{a \dots b}) c^a \cdots c^b + u^d f_{a \dots b} \partial_d (c^a \cdots c^b).$$

Given a vector bundle $E \rightarrow X$, let us consider the simple graded manifold $(J^r Y, \mathfrak{A}_{E_r})$ whose body is $J^r Y$ and the structure bundle is the pull-back

$$E_r = J^r Y \times_X E$$

onto $J^r Y$ of the jet bundle $J^r E \rightarrow X$, which is a vector bundle. There is an epimorphism of graded manifolds

$$(J^{r+1}Y, \mathfrak{A}_{E_{r+1}}) \rightarrow (J^r Y, \mathfrak{A}_{E_r}).$$

It consists of the open surjection π_r^{r+1} and the sheaf monomorphism $\pi_r^{r+1*} : \mathfrak{A}_{E_r} \rightarrow \mathfrak{A}_{E_{r+1}}$, where $\pi_r^{r+1*}\mathfrak{A}_{E_r}$ is the pull-back onto $J^{r+1}Y$ of the topological fiber bundle $\mathfrak{A}_{E_r} \rightarrow J^r Y$. These sheaf monomorphisms yield monomorphisms of the corresponding canonical presheaves $\overline{\mathfrak{A}}_{E_r} \rightarrow \overline{\mathfrak{A}}_{E_{r+1}}$, which make up a direct system

$$\overline{\mathfrak{A}}_{Y \times E} \longrightarrow \overline{\mathfrak{A}}_{E_1} \longrightarrow \cdots \overline{\mathfrak{A}}_{E_r} \longrightarrow \cdots, \quad (4)$$

and the monomorphisms of graded commutative rings $\mathcal{A}_{E_r} \rightarrow \mathcal{A}_{E_{r+1}}$ assembled into the direct system

$$\mathcal{A}_{Y \times E} \longrightarrow \mathcal{A}_{E_1} \longrightarrow \cdots \mathcal{A}_{E_r} \longrightarrow \cdots. \quad (5)$$

A direct limit of this direct system is a graded commutative ring $\mathcal{S}_\infty[E; Y]$ of all graded functions $f \in \mathcal{A}_{E_r}$ on jet manifolds $J^r Y$ modulo monomorphisms π_r^{r+1*} . The monomorphisms $C^\infty(J^r Y) \rightarrow \mathcal{A}_{E_r}$ provide the monomorphism $\mathcal{O}_\infty Y \rightarrow \mathcal{S}_\infty[E; Y]$, while the body epimorphisms $\mathcal{A}_{E_r} \rightarrow C^\infty(J^r Y)$ yield the epimorphism $\mathcal{S}_\infty[E; Y] \rightarrow \mathcal{O}_\infty Y$.

One can think of elements of $\mathcal{S}_\infty[E; Y]$ as being graded functions on $J^\infty Y$ as follows. A direct limit of the direct system of presheaves (4) is a presheaf on the infinite order jet manifold $J^\infty Y$. Let $\mathfrak{T}_\infty[E; Y]$ be the sheaf of germs of this presheaf. The structure module $\Gamma(\mathfrak{T}_\infty[E; Y])$ of sections of $\mathfrak{T}_\infty[E; Y]$ is a ring such that, given an element $f \in \Gamma(\mathfrak{T}_\infty[E; Y])$ and a point $z \in J^\infty Y$, there exist an open neighbourhood U of z and a graded function $f^{(k)}$ on some finite order jet manifold $J^k Y$ so that $f|_U = \pi_k^{\infty*} f^{(k)}|_U$. In particular, there is the monomorphism $\mathcal{S}_\infty[E; Y] \rightarrow \Gamma(\mathfrak{T}_\infty[E; Y])$. Due to this monomorphism, one can restrict $\mathcal{S}_\infty[E; Y]$ to the coordinate chart (2), and say that $\mathcal{S}_\infty[E; Y]$ as an $\mathcal{O}_\infty Y$ -ring is locally generated by the elements c_Λ^a , where $\{c_\Lambda^a\}$ is a local fiber basis for E^* over $\pi_\infty(U)$. We agree to call $(y_\Lambda^i, c_\Lambda^a)$, $0 \leq |\Lambda|$, the local basis for $\mathcal{S}_\infty[E; Y]$. Let the collective symbol s_Λ^A stand for its elements.

Remark 2. One can think of c_Λ^a as being jets of graded functions c^a [2, 5]. This definition differs from the notion of jets of a graded commutative ring [7] and that of jets of a graded fiber bundle [9], but reproduces the heuristic notion of jets of odd fields in Lagrangian field theory [10].

Remark 3. Let V , V' and V'' be vector bundles over X . Let us consider the ring

$$\mathcal{S}_\infty[V \times_X E; Y \times_X V' \times_X V''],$$

and its subring of graded functions which are polynomial in fiber coordinates of the vector bundle $Y \times_X V' \times_X V'' \rightarrow Y$. We denote the latter by $\mathcal{P}_\infty[VE; Y; V'V'']$. One can think of its elements as being graded functions on $J^\infty Y$, too.

Let $\vartheta \in \mathfrak{d}\mathcal{S}_\infty[E; Y]$ be a graded derivation of the $\mathbb{R}$ -ring $\mathcal{S}_\infty[E; Y]$ [2, 5]. With respect to the local basis (s_Λ^A) , it takes the form

$$\vartheta = \vartheta^\lambda d_\lambda + (\vartheta^A \partial_A + \sum_{|\Lambda| > 0} d_\Lambda \vartheta^A \partial_A^\Lambda),$$

where the tuple of graded derivations $\{\partial_a^\Lambda\}$ is the dual of the tuple $\{c_\Lambda^a\}$ of generating elements of the $\mathcal{O}_\infty[F; Y]$ -algebra $\mathcal{S}_\infty[E; Y]$, and $\vartheta^\lambda, \vartheta^A$ are local graded functions. We further restrict our consideration to vertical contact graded derivations

$$\vartheta = \sum_{0 \leq |\Lambda|} d_\Lambda \vartheta^A \partial_A^\Lambda. \quad (6)$$

Such a derivation is completely determined by its first summand

$$v = v^A(x^\lambda, s_\Lambda^A) \partial_A, \quad 0 \leq |\Lambda| \leq k, \quad (7)$$

called a generalized graded vector field. For the sake of simplicity, the common symbol v further stands both for a contact graded derivation (6) and a generalized vector field (7), which is also called a graded derivation. A graded derivation v is said to be nilpotent if $v(v(f)) = 0$ for any graded function $f \in \mathcal{S}_\infty[E; Y]$. One can show that v is nilpotent only if it is odd and iff all v^A obey the equality

$$v(v^A) = \sum_{0 \leq |\Sigma|} v_\Sigma^B \partial_B^\Sigma(v^A) = 0. \quad (8)$$

3 Noether identities

Let $\mathcal{E}$ be a E -valued differential operator on a smooth fiber bundle $Y \rightarrow X$.

Proposition 2. *One can associate to $\mathcal{E}$ a chain complex whose boundaries vanish on $\text{Ker } \mathcal{E}$.*

Proof. A differential operator $\mathcal{E}$ is locally represented by a set of functions $\mathcal{E}^a \in \mathcal{O}_\infty Y$, possessing the corresponding coordinate transformation law. Let us provide the ring $\mathcal{S}_\infty[E; Y]$ with the nilpotent graded derivation

$$\delta = \overleftarrow{\partial}_a \mathcal{E}^a, \quad (9)$$

whose definition is independent of the choice of a local basis. It is convenient to deal with a graded derivation δ (9) acting on graded functions on the right by the rule

$$\delta(f \wedge f') = (-1)^{[f']} \delta(f) \wedge f' + f \wedge \delta(f'),$$

where $[f']$ denotes the Grassmann parity. We call δ (9) the Koszul–Tate differential. With this differential, the ring $\mathcal{S}_\infty[E; Y]$ is split into the chain complex

$$0 \leftarrow \mathcal{O}_\infty Y \xleftarrow{\delta} \mathcal{S}_\infty[E; Y]_1 \cdots \xleftarrow{\delta} \mathcal{S}_\infty[E; Y]_k \cdots \quad (10)$$

graded in polynomials of the odd elements c_Λ^a . Following the physical literature [2, 10], we assign to c_Λ^a the antifield number $\text{Ant}[c_\Lambda^a] = 1$. It is readily observed that the boundaries of the complex (10) vanish on $\text{Ker } \mathcal{E}$. $\square$

Note that homology groups $H_*(\delta)$ of the complex (10) are $\mathcal{O}_\infty Y$ -modules, but these modules fail to be torsion-free. Indeed, given a cycle $\Phi \in \mathcal{S}_\infty[E; Y]_k$ and an element $f = \delta\sigma$ of the ring $\mathcal{O}_\infty Y \subset \mathcal{S}_\infty[E; Y]$, we obtain that $f\Phi = \delta(\sigma\Phi)$ is a boundary. Therefore, one can not apply the Künneth formula to the homology of this complex, though any its term $\mathcal{S}_\infty[E; Y]_k$ is isomorphic to the graded commutative k -tensor product of the $\mathcal{O}_\infty Y$ -module $\mathcal{S}_\infty[E; Y]_1$.

The homology $H_0(\delta)$ of the complex (10) is not trivial, but this homology and the higher ones $H_{k \geq 2}(\delta)$ are not essential for our consideration. Therefore, we replace the complex (10) with the finite one

$$0 \leftarrow \text{Im } \delta \xleftarrow{\delta} \mathcal{S}_\infty[E; Y]_1 \xleftarrow{\delta} \mathcal{S}_\infty[E; Y]_2 \quad (11)$$

of graded functions of antifield number $k \leq 2$. It is exact at $\text{Im } \delta$, and its first homology coincides with that of the complex (10). Let us consider this homology.

A generic one-chain of the complex (11) takes the form

$$\Phi = \sum_{0 \leq |\Lambda|} \Phi_a^\Lambda c_\Lambda^a, \quad \Phi_a^\Lambda \in \mathcal{O}_\infty Y, \quad (12)$$

and the cycle condition $\delta\Phi = 0$ reads

$$\sum_{0 \leq |\Lambda|} \Phi_a^\Lambda d_\Lambda \mathcal{E}^a = 0. \quad (13)$$

One can think of this equality as being a reduction condition on a differential operator $\mathcal{E}$. Conversely, any reduction condition of form (13) comes from some cycle (12). The reduction condition (13) is trivial if a cycle is a boundary, i.e., it takes the form

$$\Phi = \sum_{0 \leq |\Lambda|, |\Sigma|} T_{ab}^{\Lambda\Sigma} d_\Sigma \mathcal{E}^b c_\Lambda^a, \quad T_{ab}^{\Lambda\Sigma} = -T_{ba}^{\Sigma\Lambda}. \quad (14)$$

If $\mathcal{E}$ is an Euler–Lagrange operator of some Lagrangian system on a fiber bundle $Y \rightarrow X$, the nontrivial reduction condition (13) is a Noether identity [1, 2]. Therefore, we come to the following definition.

Definition 3. *A differential operator $\mathcal{E}$ is called degenerate if the homology $H_1(\delta)$ of the complex (14) (or (12)) is not trivial. We agree to call a cycle condition (13) the Noether identity.*

One can say something more if the $\mathcal{O}_\infty Y$ -module $H_1(\delta)$ is finitely generated, i.e., it possesses the following particular structure. There are elements $\Delta \in H_1(\delta)$ making up a projective $C^\infty(X)$ -module $\mathcal{C}_{(0)}$ of finite rank which, by virtue of the Serre–Swan theorem,

is isomorphic to the module of sections of some vector bundle $E_0 \rightarrow X$. Let $\{\Delta^r\}$ be local bases for this $C^\infty(X)$ -module. Then every element $\Phi \in H_1(\delta)$ factorizes

$$\Phi = \sum_{0 \leq |\Xi|} G_r^\Xi d_\Xi \Delta^r, \quad G_r^\Xi \in \mathcal{O}_\infty Y, \quad (15)$$

$$\Delta^r = \sum_{0 \leq |\Lambda|} \Delta_a^{\Lambda r} c_\Lambda^a, \quad \Delta_a^{\Lambda r} \in \mathcal{O}_\infty Y, \quad (16)$$

via elements of $\mathcal{C}_{(0)}$, i.e., any Noether identity (13) is a corollary of the Noether identities

$$\sum_{0 \leq |\Lambda|} \Delta_a^{\Lambda r} d_\Lambda \mathcal{E}^a = 0. \quad (17)$$

Clearly, the factorization (15) is independent of specification of local bases $\{\Delta^r\}$. We say that the Noether identities (17) are complete, and call $\Delta \in \mathcal{C}_{(0)}$ the Noether operators. Note that, being representatives of $H_1(\delta)$, the graded functions Δ^r (16) are not δ -exact.

Proposition 4. *If the homology $H_1(\delta)$ of the complex (11) is finitely generated, this complex can be extended to a one-exact complex with a boundary operator whose nilpotency conditions (8) are just complete Noether identities (see the complex (19) below).*

Proof. Let us consider the ring $\mathcal{P}_\infty[E; Y; E_0]$ of graded functions on $J^\infty Y$ (see Remark 3). It possesses local bases $\{y_\Lambda^i, c_\Lambda^a, c_\Lambda^r\}$, where $[c_\Lambda^r] = 0$ and $\text{Ant}[c_\Lambda^r] = 2$. This ring is provided with the nilpotent graded derivation

$$\delta_0 = \delta + \overleftarrow{\partial}_r \Delta^r, \quad (18)$$

called the extended Koszul–Tate differential. Its nilpotency conditions (8) are equivalent to the complete Noether identities (17). Then the module $\mathcal{P}_\infty[E; Y; E_0]_{\leq 3}$ of graded functions of antifield number $k \leq 3$ is split into the chain complex

$$0 \leftarrow \text{Im } \delta \xleftarrow{\delta} \mathcal{S}_\infty[E; Y]_1 \xleftarrow{\delta_0} \mathcal{P}_\infty[E; Y; E_0]_2 \xleftarrow{\delta_0} \mathcal{P}_\infty[E; Y; E_0]_3. \quad (19)$$

Let $H_*(\delta_0)$ denote its homology. We have $H_0(\delta_0) = H_0(\delta) = 0$. Furthermore, any one-cycle Φ up to a boundary takes the form (15) and, therefore, it is a δ_0 -boundary

$$\Phi = \sum_{0 \leq |\Sigma|} G_r^\Sigma d_\Sigma \Delta^r = \delta_0 \left(\sum_{0 \leq |\Sigma|} G_r^\Sigma c_\Sigma^r \right).$$

Hence, $H_1(\delta_0) = 0$, i.e., the complex (19) is one-exact. $\square$

4 The Koszul–Tate complex

Turn now to the second homology $H_2(\delta_0)$ of the complex (19). A generic two-chain reads

$$\Phi = G + H = \sum_{0 \leq |\Lambda|} G_r^\Lambda c_\Lambda^r + \sum_{0 \leq |\Lambda|, |\Sigma|} H_{ab}^{\Lambda\Sigma} c_\Lambda^a c_\Sigma^b, \quad G_r^\Lambda, H_{ab}^{\Lambda\Sigma} \in \mathcal{O}_\infty Y. \quad (20)$$

The cycle condition $\delta_0 \Phi = 0$ takes the form

$$\sum_{0 \leq |\Lambda|} G_r^\Lambda d_\Lambda \Delta^r + \delta H = 0. \quad (21)$$

One can think of this equality as being the reduction condition on the Noether operators Δ^r (16). Conversely, let

$$\Phi = \sum_{0 \leq |\Lambda|} G_r^\Lambda c_\Lambda^r \in \mathcal{P}_\infty[E; Y; E_0]_2$$

be a graded function such that the reduction condition (21) holds. Obviously, it is a cycle condition of the two-chain (20). The reduction condition (21) is trivial either if a two-cycle Φ (20) is a boundary or its summand G vanishes on $\text{Ker } \mathcal{E}$.

Definition 5. *A degenerate differential operator is said to be one-stage reducible if there exist non-trivial reduction conditions (21), called first-stage Noether identities.*

Proposition 6. *First-stage Noether identities can be identified to nontrivial elements of the homology $H_2(\delta_0)$ iff any δ -cycle $\Phi \in \mathcal{S}_\infty[E; Y]_2$ is a δ_0 -boundary.*

Proof. It suffices to show that, if the summand G of a two-cycle Φ (20) is δ -exact, then Φ is a boundary. If $G = \delta\Psi$, then

$$\Phi = \delta_0 \Psi + (\delta - \delta_0) \Psi + H. \quad (22)$$

The cycle condition reads

$$\delta_0 \Phi = \delta((\delta - \delta_0) \Psi + H) = 0.$$

Then $(\delta - \delta_0) \Psi + H$ is δ_0 -exact since any δ -cycle $\phi \in \mathcal{S}_\infty[E; Y]_2$ by assumption is a δ_0 -boundary. Consequently, Φ (22) is δ_0 -exact. Conversely, let $\Phi \in \mathcal{S}_\infty[E; Y]_2$ be an arbitrary δ -cycle. The cycle condition reads

$$\delta \Phi = 2\Phi_{ab}^{\Lambda\Sigma} c_\Lambda^a \delta c_\Sigma^b = 2\Phi_{ab}^{\Lambda\Sigma} c_\Lambda^a d_\Sigma \mathcal{E}^b = 0. \quad (23)$$

It follows that $\Phi_{ab}^{\Lambda\Sigma} \delta c_\Sigma^b = 0$ for all induces (a, Λ) . Omitting a δ -boundary term, we obtain

$$\Phi_{ab}^{\Lambda\Sigma} c_\Sigma^b = G_{ar}^{\Lambda\Xi} d_\Xi \Delta^r.$$

Hence, Φ takes the form

$$\Phi = -G_{ar}^{\Lambda\Xi} d_\Xi \Delta^r c_\Lambda^a. \quad (24)$$

We can associate to it the three-chain $\Psi = -G_{ar}^{\Lambda\Xi} c_\Xi^r c_\Lambda^a$ such that

$$\delta_0 \Psi = \Phi + \sigma = \Phi - G_{ar}^{\Lambda\Xi} d_\Lambda \mathcal{E}^a c_\Xi^r.$$

Owing to the equality $\delta \Phi = 0$, we have $\delta_0 \sigma = 0$. Since σ is δ -exact, it by assumption is δ_0 -exact, i.e., $\sigma = \delta_0 \psi$. Then we obtain that $\Phi = \delta_0 \Psi - \delta_0 \psi$. $\square$

Lemma 7. *It is easily justified that a two-cycle $\Phi \in \mathcal{S}_\infty[E; Y]_2$ is δ_0 -exact iff Φ up to a δ -boundary takes the form*

$$\Phi = \sum_{0 \leq |\Lambda|, |\Sigma|} G_{rr'}^{\Sigma\Lambda} d_\Sigma \Delta^r d_\Lambda \Delta^{r'}. \quad (25)$$

If the condition of Proposition 6 (called the two-homology regularity condition) is satisfied, let us assume that first-stage Noether identities are finitely generated as follows. There are elements $\Delta_{(1)} \in H_2(\delta_0)$ making up a projective $C^\infty(X)$ -module $\mathcal{C}_{(1)}$ of finite rank which is isomorphic to the module of sections of some vector bundle $E_1 \rightarrow X$. Let $\{\Delta^{r_1}\}$ be local bases for this $C^\infty(X)$ -module. Every element $\Phi \in H_2(\delta_0)$ factorizes

$$\Phi = \sum_{0 \leq |\Xi|} \Phi_{r_1}^\Xi d_\Xi \Delta^{r_1}, \quad \Phi_{r_1}^\Xi \in \mathcal{O}_\infty Y, \quad (26)$$

$$\Delta^{r_1} = G^{r_1} + h^{r_1} = \sum_{0 \leq |\Lambda|} \Delta_r^{\Lambda r_1} c_\Lambda^r + h^{r_1}, \quad h^{r_1} \in \mathcal{S}_\infty[E; Y]_2, \quad (27)$$

via elements of $\mathcal{C}_{(1)}$, i.e., any first-stage Noether identity (21) results from the equalities

$$\sum_{0 \leq |\Lambda|} \Delta_r^{\Lambda r_1} d_\Lambda \Delta^r + \delta h^{r_1} = 0, \quad (28)$$

called the complete first-stage Noether identities. Elements of $\mathcal{C}_{(1)}$ are said to be the first-stage Noether operators. Note that the first summands G^{r_1} of the operators Δ^{r_1} (27) are not δ -exact.

Proposition 8. *Given a reducible degenerate differential operator $\mathcal{E}$, let the associated one-exact complex (19) obey the two-homology regularity condition and let its homology $H_2(\delta_0)$ be finitely generated. Then this complex is extended to the two-exact one with a boundary operator whose nilpotency conditions are equivalent to complete Noether and first-stage Noether identities (see the complex (30) below).*

Proof. Let us consider the ring $\mathcal{P}_\infty^*[E_1 E; Y; E_0]$ of graded functions on $J^\infty Y$ possessing local bases $\{y_\Lambda^i, c_\Lambda^a, c_\Lambda^r, c_\Lambda^{r_1}\}$, where $[c_\Lambda^{r_1}] = 1$ and $\text{Ant}[c_\Lambda^{r_1}] = 3$. It can be provided the first-stage Koszul–Tate differential defined as the nilpotent graded derivation

$$\delta_1 = \delta_0 + \overleftarrow{\partial}_{r_1} \Delta^{r_1}. \quad (29)$$

Its nilpotency conditions (8) are equivalent to complete Noether identities (17) and complete first-stage Noether identities (28). Then the module $\mathcal{P}_\infty[E_1 E; Y; E_0]_{\leq 4}$ of graded functions of antifield number $\text{Ant}[\phi] \leq 4$ is split into the chain complex

$$\begin{aligned} 0 \leftarrow \text{Im } \delta \xleftarrow{\delta} \mathcal{S}_\infty[E; Y]_1 \xleftarrow{\delta_0} \mathcal{P}_\infty[E; Y; E_0]_2 \xleftarrow{\delta_1} \mathcal{P}_\infty[E_1 E; Y; E_0]_3 \\ \xleftarrow{\delta_1} \mathcal{P}_\infty[E_1 E; Y; E_0]_4. \end{aligned} \quad (30)$$

Let $H_*(\delta_1)$ denote its homology. It is readily observed that

$$H_0(\delta_1) = H_0(\delta) = 0, \quad H_1(\delta_1) = H_1(\delta_0) = 0.$$

By virtue of the expression (26), any two-cycle of the complex (30) is a boundary

$$\Phi = \sum_{0 \leq |\Xi|} \Phi_{r_1}^\Xi d_\Xi \Delta^{r_1} = \delta_1 \left(\sum_{0 \leq |\Xi|} \Phi_{r_1}^\Xi c_\Xi^{r_1} \right).$$

It follows that $H_2(\delta_1) = 0$, i.e., the complex (30) is two-exact. $\square$

If the third homology $H_3(\delta_1)$ of the complex (30) is not trivial, there are reduction conditions on the first-stage Noether operators, and so on. Iterating the arguments, we come to the following.

Let $\mathcal{E}$ be a degenerate differential operator whose Noether identities are finitely generated. In accordance with Proposition 4, we associate to it the one-exact chain complex (19). Given an integer $N \geq 1$, let $E_1, \dots, E_N$ be some vector bundles over X . Let us consider the ring

$$\mathcal{P}_\infty\{N\} = \mathcal{P}_\infty[E_{N-1} \cdots E_1 E; Y; E_0 \cdots E_N]$$

of graded functions on $J^\infty Y$ if N is even or the ring

$$\mathcal{P}_\infty\{N\} = \mathcal{P}_\infty[E_N \cdots E_1 E; Y; E_0 \cdots E_{N-1}]$$

if N is odd. It possesses local bases $\{y_\Lambda^i, c_\Lambda^a, c_\Lambda^r, c_\Lambda^{r_1}, \dots, c_\Lambda^{r_N}\}$ where $[c_\Lambda^{r_k}] = (k+1) \bmod 2$ and $\text{Ant}[c_\Lambda^{r_k}] = k+2$. Let $k = -1, 0$ further stand for y^i and c^r , respectively. We assume that:

(i) the ring $\mathcal{P}_\infty\{N\}$ is provided with the nilpotent graded derivation

$$\delta_N = \delta_0 + \sum_{1 \leq k \leq N} \overset{\leftarrow}{\partial}_{r_k} \Delta^{r_k}, \quad (31)$$

$$\Delta^{r_k} = G^{r_k} + h^{r_k} = \sum_{0 \leq |\Lambda|} \Delta_{r_{k-1}}^{\Lambda r_k} c_\Lambda^{r_{k-1}} + \sum_{0 \leq \Sigma, 0 \leq \Xi} (h_{ar_{k-2}}^{\Xi \Sigma r_k} c_\Xi^a c_\Sigma^{r_{k-2}} + \dots), \quad (32)$$

of antifield number -1;

(ii) the module $\mathcal{P}_\infty\{N\}_{\leq N+3}$ of graded functions of antifield number $k \leq N+3$ is split into the $(N+1)$ -exact chain complex

$$\begin{aligned} 0 \leftarrow \text{Im } \delta \overset{\delta}{\longleftarrow} \mathcal{S}_\infty[E; Y]_1 \overset{\delta_0}{\longleftarrow} \mathcal{P}_\infty\{0\}_2 \overset{\delta_1}{\longleftarrow} \mathcal{P}_\infty\{1\}_3 \cdots \\ \overset{\delta_{N-1}}{\longleftarrow} \mathcal{P}_\infty\{N-1\}_{N+1} \overset{\delta_N}{\longleftarrow} \mathcal{P}_\infty\{N\}_{N+2} \overset{\delta_N}{\longleftarrow} \mathcal{P}_\infty\{N\}_{N+3}, \end{aligned} \quad (33)$$

which satisfies the $(N+1)$ -homology regularity condition, introduced below.

Definition 9. One says that the complex (33) obeys the $(N+1)$ -homology regularity condition if any $\delta_{k < N-1}$ -cycle $\Phi \in \mathcal{P}_\infty\{k\}_{k+3} \subset \mathcal{P}_\infty\{k+1\}_{k+3}$ is a δ_{k+1} -boundary.

Remark 4. The $(N+1)$ -exactness of the complex (33) implies that any $\delta_{k < N-1}$ -cycle $\Phi \in \mathcal{P}_\infty^{0,n}\{k\}_{k+3}$, $k < N$, is a δ_{k+2} -boundary, but not necessary a δ_{k+1} -one.

If $N = 1$, the complex $\mathcal{P}_\infty\{1\}_{\leq 4}$ (33) restarts the complex (30) associated to a first-stage reducible differential operator in accordance with Proposition 8. Therefore, we agree to call

δ_N (31) the N -stage Koszul–Tate differential. Its nilpotency implies complete Noether identities (17), first-stage Noether identities (28) and the equalities

$$\sum_{0 \leq |\Lambda|} \Delta_{r_{k-1}}^{\Lambda} d_{\Lambda} \left(\sum_{0 \leq |\Sigma|} \Delta_{r_{k-2}}^{\Sigma} c_{\Sigma}^{r_{k-2}} \right) + \delta \left(\sum_{0 \leq \Sigma, 0 \leq \Xi} h_{ar_{k-2}}^{\Xi \Sigma} c_{\Xi}^a c_{\Sigma}^{r_{k-2}} \right) = 0, \quad (34)$$

for $k = 2, \dots, N$. One can think of the equalities (34) as being complete k -stage Noether identities because of their properties which we will justify in the case of $k = N + 1$. Accordingly, Δ^{r_k} (32) are said to be the k -stage Noether operators.

A generic $(N + 2)$ -chain $\Phi \in \mathcal{P}_{\infty}\{N\}_{N+2}$ takes the form

$$\Phi = G + H = \sum_{0 \leq |\Lambda|} G_{r_N}^{\Lambda} c_{\Lambda}^{r_N} + \sum_{0 \leq \Sigma, 0 \leq \Xi} (H_{ar_{N-1}}^{\Xi \Sigma} c_{\Xi}^a c_{\Sigma}^{r_{N-1}} + \dots). \quad (35)$$

Let it be a cycle. The cycle condition $\delta_N \Phi = 0$ implies the equality

$$\sum_{0 \leq |\Lambda|} G_{r_N}^{\Lambda} d_{\Lambda} \left(\sum_{0 \leq |\Sigma|} \Delta_{r_{N-1}}^{\Sigma} c_{\Sigma}^{r_{N-1}} \right) + \delta \left(\sum_{0 \leq \Sigma, 0 \leq \Xi} H_{ar_{N-1}}^{\Xi \Sigma} c_{\Xi}^a c_{\Sigma}^{r_{N-1}} \right) = 0. \quad (36)$$

One can think of this equality as being the reduction condition on the N -stage Noether operators (32). Conversely, let

$$\Phi = \sum_{0 \leq |\Lambda|} G_{r_N}^{\Lambda} c_{\Lambda}^{r_N} \in \mathcal{P}_{\infty}\{N\}_{N+2}$$

be a graded function such that the reduction condition (36) holds. Then this reduction condition can be extended to a cycle one as follows. It is brought into the form

$$\begin{aligned} \delta_N \left(\sum_{0 \leq |\Lambda|} G_{r_N}^{\Lambda} c_{\Lambda}^{r_N} + \sum_{0 \leq \Sigma, 0 \leq \Xi} H_{ar_{N-1}}^{\Xi \Sigma} c_{\Xi}^a c_{\Sigma}^{r_{N-1}} \right) = \\ - \sum_{0 \leq |\Lambda|} G_{r_N}^{\Lambda} d_{\Lambda} h^{r_N} + \sum_{0 \leq \Sigma, 0 \leq \Xi} H_{ar_{N-1}}^{\Xi \Sigma} c_{\Xi}^a d_{\Sigma} \Delta^{r_{N-1}}. \end{aligned}$$

A glance at the expression (32) shows that the term in the right-hand side of this equality belongs to $\mathcal{P}_{\infty}\{N - 2\}_{N+1}$. It is a δ_{N-2} -cycle and, consequently, a δ_{N-1} -boundary $\delta_{N-1} \Psi$ in accordance with the $(N + 1)$ -homology regularity condition. Then the reduction condition (36) is a $c_{\Sigma}^{r_{N-1}}$ -dependent part of the cycle condition

$$\delta_N \left(\sum_{0 \leq |\Lambda|} G_{r_N}^{\Lambda} c_{\Lambda}^{r_N} + \sum_{0 \leq \Sigma, 0 \leq \Xi} H_{ar_{N-1}}^{\Xi \Sigma} c_{\Xi}^a c_{\Sigma}^{r_{N-1}} - \Psi \right) = 0,$$

but $\delta_N \Psi$ does not make a contribution to this reduction condition.

Being a cycle condition, the reduction condition (36) is trivial either if a cycle Φ (35) is a δ_N -boundary or its summand G is δ -exact, i.e., it is a boundary, too, as we have stated above. Then Definition 5 can be generalized as follows.

Definition 10. A degenerate differential operator is said to be $(N + 1)$ -stage reducible if there exist non-trivial reduction conditions (36), called the $(N + 1)$ -stage Noether identities.

Theorem 11. (i) The $(N + 1)$ -stage Noether identities can be identified to nontrivial elements of the homology $H_{N+2}(\delta_N)$ of the complex (33) iff this homology obeys the $(N + 2)$ -homology regularity condition. (ii) If the homology $H_{N+2}(\delta_N)$ is finitely generated as defined below, the complex (33) admits an $(N + 2)$ -exact extension.

Proof. (i) The $(N + 2)$ -homology regularity condition implies that any δ_{N-1} -cycle $\Phi \in \mathcal{P}_\infty\{N - 1\}_{N+2} \subset \mathcal{P}_\infty\{N\}_{N+2}$ is a δ_N -boundary. Therefore, if Φ (35) is a representative of a nontrivial element of $H_{N+2}(\delta_N)$, its summand G linear in $c_\Lambda^{r_N}$ does not vanish. Moreover, it is not a δ -boundary. Indeed, if $\Phi = \delta\Psi$, then

$$\Phi = \delta_N\Psi + (\delta - \delta_N)\Psi + H. \quad (37)$$

The cycle condition takes the form

$$\delta_N\Phi = \delta_{N-1}((\delta - \delta_N)\Psi + H) = 0.$$

Hence, $(\delta - \delta_0)\Psi + H$ is δ_N -exact since any δ_{N-1} -cycle $\phi \in \mathcal{P}_\infty\{N - 1\}_{N+2}$ is a δ_N -boundary. Consequently, Φ (37) is a boundary. If the $(N + 2)$ -homology regularity condition does not hold, trivial reduction conditions (36) also come from nontrivial elements of the homology $H_{N+2}(\delta_N)$.

(ii) Let the $(N + 1)$ -stage Noether identities be finitely generated. Namely, there exist elements $\Delta_{(N+1)} \in H_{N+2}(\delta_N)$ making up a projective $C^\infty(X)$ -module $\mathcal{C}_{(N+1)}$ of finite rank which is isomorphic to the module of sections of some vector bundle $E_{N+1} \rightarrow X$. Let $\{\Delta^{r_{N+1}}\}$ be local bases for this $C^\infty(X)$ -module. Then any element $\Phi \in H_{N+2}(\delta_N)$ factorizes

$$\Phi = \sum_{0 \leq |\Xi|} \Phi_{r_{N+1}}^\Xi d_\Xi \Delta^{r_{N+1}}, \quad \Phi_{r_{N+1}}^\Xi \in \mathcal{O}_\infty Y, \quad (38)$$

$$\Delta^{r_{N+1}} = G^{r_{N+1}} + h^{r_{N+1}} = \sum_{0 \leq |\Lambda|} \Delta_{r_N}^{\Lambda r_{N+1}} c_\Lambda^{r_N} + h^{r_{N+1}}, \quad (39)$$

through elements of $\mathcal{C}_{(N+1)}$. Clearly, this factorization is independent of specification of local bases $\{\Delta^{r_{N+1}}\}$. Let us extend the ring $\mathcal{P}_\infty\{N\}$ to the ring $\mathcal{P}_\infty\{N + 1\}$ possessing local bases

$$\{y_\Lambda^i, c_\Lambda^a, c_\Lambda^r, c_\Lambda^{r_1}, \dots, c_\Lambda^{r_N}, c_\Lambda^{r_{N+1}}\}, \quad [c_\Lambda^{r_{N+1}}] = N \bmod 2, \quad \text{Ant}[c_\Lambda^{r_{N+1}}] = N + 3.$$

It is provided with the nilpotent graded derivation

$$\delta_{N+1} = \delta_N + \overleftarrow{\partial}_{r_{N+1}} \Delta^{r_{N+1}} \quad (40)$$

of antifield number -1. With this graded derivation, the module $\mathcal{P}_\infty\{N+1\}_{\leq N+4}$ of graded functions of antifield number $\text{Ant}[f] \leq N+4$ is split into the chain complex

$$\begin{aligned} 0 \leftarrow \text{Im } \delta \xleftarrow{\delta} \mathcal{S}_\infty[E; Y]_1 \xleftarrow{\delta_0} \mathcal{P}_\infty\{0\}_2 \xleftarrow{\delta_1} \mathcal{P}_\infty\{1\}_3 \cdots \xleftarrow{\delta_{N-1}} \mathcal{P}_\infty\{N-1\}_{N+1} \\ \xleftarrow{\delta_N} \mathcal{P}_\infty\{N\}_{N+2} \xleftarrow{\delta_{N+1}} \mathcal{P}_\infty\{N+1\}_{N+3} \xleftarrow{\delta_{N+1}} \mathcal{P}_\infty\{N+1\}_{N+4}. \end{aligned} \quad (41)$$

It is readily observed that this complex is $(N+2)$ -exact. In this case, the $(N+1)$ -stage Noether identities (36) come from the complete $(N+1)$ -stage Noether identities

$$\sum_{0 \leq |\Lambda|} \Delta_{r_N}^{\Lambda r_{N+1}} d_\Lambda \Delta^{r_N} + \delta h^{r_{N+1}} = 0,$$

which are reproduced as the nilpotency conditions of the graded derivation (40). $\square$

The iteration procedure based on Theorem 11 can be prolonged up to an integer $N_{\max}$ when the $N_{\max}$ -stage Noether identities are irreducible, i.e., the homology $H_{N_{\max}+2}(\delta_{N_{\max}})$ is trivial. This iteration procedure may also be infinite. It results in the manifested exact Koszul–Tate complex with the Koszul–Tate boundary operator whose nilpotency conditions reproduce all Noether and higher Noether identities of an original differential operator $\mathcal{E}$.

References

- [1] D. Bashkirov, G. Giachetta, L. Mangiarotti, and G. Sardanashvily, Noether’s second theorem in a general setting. Reducible gauge theories, *J. Phys. A* **38** (2005), 5239–5344.
- [2] D. Bashkirov, G. Giachetta, L. Mangiarotti, and G. Sardanashvily, Noether’s second theorem for BRST symmetries, *J. Math. Phys.* **46** (2005), 053517.
- [3] I.Krasil’shchik, V.Lychagin and A.Vinogradov, *Geometry of Jet Spaces and Nonlinear Partial Differential Equations* (Gordon and Breach, Glasgow, 1985).
- [4] R.Bryant, S.Chern, R.Gardner, H.Goldschmidt and P.Griffiths, *Exterior Differential Systems* (Springer-Verlag, Berlin, 1991).
- [5] G. Giachetta, L. Mangiarotti, and G. Sardanashvily, Lagrangian supersymmetries depending on derivatives. Global analysis and cohomology, *Commun. Math. Phys.* (2005) (accepted).
- [6] C. Bartocci, U. Bruzzo, and D. Hernández Ruipérez, *The Geometry of Supermanifolds* (Kluwer, Dordrecht, 1991).
- [7] G. Giachetta, L. Mangiarotti, and G. Sardanashvily, *Geometric and Algebraic Topological Methods in Quantum Mechanics* (World Scientific, Singapore, 2005).

- [8] A. Rennie, Smoothnes and locality for nonunital spectral triples, *K-Theory* **28** (2003), 127-165.
- [9] D. Hernández Ruipérez and J. Muñoz Masqué, Global variational calculus on graded manifolds, *J. Math. Pures Appl.* **63** (1984), 283-309.
- [10] G. Barnich, F. Brandt and M. Henneaux, Local BRST cohomology in gauge theories, *Phys. Rep.* **338** (2000), 439-569.